



Transferability and characterization of microsatellite markers in two Neotropical *Ficus* species

Alison Gonçalves Nazareno¹, Rodrigo Augusto Santinelo Pereira², Juliana Massimino Feres³, Moacyr Antônio Mestriner³ and Ana Lilia Alzate-Marin³

¹Programa de Pós-graduação em Biologia Comparada, Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, SP, Brazil.

²Departamento de Biologia, Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, SP, Brazil.

³Laboratório de Genética Vegetal, Departamento de Genética, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, SP, Brazil.

Abstract

Microsatellite markers were transferred and characterized for two Neotropical fig tree species, *Ficus citrifolia* and *Ficus eximia*. Our study demonstrated that microsatellite markers developed from different subgenera of *Ficus* can be transferred to related species. In the present case, 12 of the 15 primer pairs tested (80%) were successfully transferred to both of the above species. Eleven loci were polymorphic when tested across 60 *F. citrifolia* and 60 *F. eximia* individuals. For *F. citrifolia*, there were 4 to 15 alleles per locus, whereas expected heterozygosities ranged from 0.31 to 0.91. In the case of *F. eximia*, this was 2 to 12 alleles per locus and expected heterozygosities from 0.42 to 0.87.

Key words: ecological genetics, *Ficus citrifolia*, *Ficus eximia*, Moraceae, SSRs.

Received: December 3, 2008; Accepted: March 23, 2009.

Effective strategies for the conservation of genetic resources in tropical forests are of great importance, mainly due to the negative impacts arising from the reduction in biological diversity. This is especially true with regard to ecologically important species such as fig trees (*Ficus* species, family Moraceae), which are considered to be keystone-resources in tropical forests, through supplying frugivores with fruit during periods of food-scarcity (Shanahan *et al.*, 2001). Furthermore, plants of the genus *Ficus* are considered as a classic example of plant-insect mutualism (Weiblen, 2002). With few exceptions, each of the 750 *Ficus* species maintains an obligatory symbiotic interaction with a specific pollinating wasp species (Hymenoptera: Agaonidae). However, little is known about the genetic diversity and population structure of *Ficus* species (Dick *et al.*, 2008). Microsatellite markers (simple sequence repeats - SSR) are informative tools used to assess the genetic structure of populations as well as basic quantitative genetic parameters. Although microsatellite markers constitute informative systems for ecological genetics, they have

only been isolated for seven of the 750 species of *Ficus* (Khadari *et al.*, 2001; Giraldo *et al.*, 2005; Zavodna *et al.*, 2005; Vignes *et al.*, 2006; Ahmed *et al.*, 2007; Bandelj *et al.*, 2007; Crozier *et al.*, 2007). Nevertheless, the high transferability of these markers has allowed for cross amplification in 47 *Ficus* species (Khadari *et al.*, 2001; Giraldo *et al.*, 2005; Vignes *et al.*, 2006). Moreover, indications of high transferability within a particular genus has also come to light from other areas of research (Poncet *et al.*, 2004; Moon *et al.*, 2008). Given the time consuming and relatively costly process of isolating microsatellites and the low frequency of SSRs in plants (Powell *et al.*, 1996), it is a decided advantage to be able to utilize primer sequences identified in one species in other closely related ones. Here, we examine the transferability and the characterization of microsatellite markers previously developed from different subgenera of *Ficus* for two species occurring in Brazil, *Ficus citrifolia* P. Miller and *Ficus eximia* Schott.

The subgenus *Urostigma* section *Americana*, to which *F. citrifolia* and *F. eximia* belong, includes monoecious plants that may occur as trees of hemi-epiphyte growth form (Berg, 1989). *Ficus citrifolia* normally grows as a hemiepiphyte on other trees or buildings and frequently develops within disturbed areas. *Ficus eximia* usually germinates on fallen trunks and grows as a free-standing tree in

humid patches in the forest. During January and February, 2008, we sampled 120 individuals from two natural populations (30 individuals per area per species), 350 km apart, located at the Parque Estadual Morro do Diabo (22° 27' - 22° 40' S, 52° 10' - 52° 22' W) and at the Estação Ecológica de Caetetus (22° 41' - 22° 46' S, 49° 10' - 49° 16' W), both in southeastern Brazil.

DNA for microsatellite analysis was extracted from frozen leaves by using the cetyltrimethyl ammonium bromide (CTAB) extraction method (Doyle and Doyle, 1990). Fifteen microsatellite loci, previously developed for *Ficus* (*Pharmacosycea*) *insipida* (Vignes *et al.*, 2006), *Ficus* (*Sycomoros*) *racemosa* and *Ficus* (*Urostigma*) *rubiginosa* (Crozier *et al.*, 2007), were tested for cross amplification in specimens of *F. citrifolia* and *F. eximia*. Using polymerase chain reaction (PCR), a screening of each primer pair through ten annealing temperatures (between 46-55 °C) was accomplished with 10 individuals of *F. citrifolia* and *F. eximia*. Microsatellite loci were amplified in a final volume of 10 µL containing 0.3 µM of each primer, 1 U *Taq* DNA polymerase, 0.25 mM of each dNTP, 1 x MgCl₂-free reaction buffer [75 mM Tris-HCl pH 9.0, 50 mM KCl and 20 mM (NH₄)₂SO₄], 1.5 mM MgCl₂ and 2.5 ng of template DNA. The amplification was performed using a Master-Cycler Eppendorf under the following conditions: 5 min of denaturation at 96 °C and 30 cycles of 30 s of initial denaturation at 94 °C, 1 min of annealing at T_a (Table 1) and 1 min of extension at 72 °C, to finish with 7 min of elongation at 72 °C. Amplified fragments were separated on 10% denaturing polyacrylamide gels in 8 M urea and 1 x TBE buffer, to then be stained with silver nitrate (Sanguinetti *et al.*, 1994). The quantification of allele size was scored against a 10 bp DNA ladder standard (Invitrogen).

Genetic diversity parameters and probabilities of paternity exclusion were estimated using CERVUS version 3.0 (Kalinowski *et al.*, 2007). The FSTAT software package version 1.2 (Goudet, 2001) was used to test all loci for linkage disequilibrium, with application of Bonferroni correction for multiple comparisons.

Our study demonstrated that microsatellite markers developed from different subgenera of *Ficus* can be transferred to related species. We successfully transferred 12 of the 15 primer pairs tested (80%) to both of the species, *F. citrifolia* and *F. eximia* (Table 1). Similar allele numbers and length of amplification products were apparent in most of the successful loci, when compared with the species from which they were developed (Table 1). Of the 12 loci transferred, 11 were polymorphic for both *F. citrifolia* and *F. eximia*. Loci FinsT7 and Frub154 were monomorphic in only one species. Eighty-seven *F. citrifolia* and 77 *F. eximia* allelic variants were identified (Table 1). Furthermore, the average for heterozygosity and the mean number of alleles per loci were, respectively, 0.67 and 7.3 in *F. citrifolia* and 0.69 and 6.4 in *F. eximia*. Heterozygosity val-

ues of *F. citrifolia* and *F. eximia* in this study were within those for *Ficus* species reported in previous studies (Bandelj *et al.*, 2007; Crozier *et al.*, 2007). The level of heterozygosity found in a population is highly dependent on the mating system and the evolutionary history of the species, besides a range of other factors. Although various microsatellite markers and sample sizes have been used in diversity studies on *Ficus* species, these same values made it possible for us to assume the present status of genetic variability due to the mating system and plant-insect mutualism. Pollen and diaspores in *Ficus* species are dispersed over long distances (Kinnaird *et al.*, 1996; Nason *et al.*, 1996), thereby implying that the flight distance of pollinators and dispersal range might predict high levels of genetic variation in these species (Hamrick and Loveless, 1989; Epperson and Alvarez-Buylla, 1997; Nazareno and Carvalho, 2008).

There were significant deviations from Hardy-Weinberg equilibrium (HWE) in seven *F. citrifolia* loci. As to *F. eximia*, nine loci were not in HWE (Table 1). Either the intrapopulation substructure produced by the sampling effect or the presence of null alleles may have caused these deviations, since analyses at the population level showed deviations from HWE for these same loci. Furthermore, as these results were obtained by using microsatellites developed in another species, the probability of a null allele occurring would be much higher than in the case of testing in the species from which they were isolated (Kim *et al.*, 2004). In future studies, the influence of null alleles on the transferability of microsatellite markers and their applicability in other species should be investigated.

The chi-square test for the independent segregation hypothesis indicated that all loci for *F. citrifolia* were in linkage equilibrium. As to *F. eximia*, however, significant linkage disequilibrium was found for the loci Frub38 and Frub415. The combined values for probabilities of paternity exclusion in all the 11 polymorphic loci were 0.996 for *F. citrifolia* and 0.995 for *F. eximia*. Using the 11 polymorphic loci enabled us to distinguish all the 60 *F. citrifolia* and 60 *F. eximia* individuals in two populations from southeastern Brazil. Hence, transferred microsatellite markers should allow for detailed parentage studies in natural populations, even in situations where both maternity and paternity are unknown. Moreover, these microsatellite loci could be highly useful for providing data on population genetics.

The high transferability of microsatellite markers developed from different subgenera of *Ficus* for *F. citrifolia* and *F. eximia* confirm the general applicability of *Ficus* microsatellite primers to this very large genus. Currently, we are using these markers to investigate the impact of tropical deforestation on population structure and genetic diversity in forest fragments in the Atlantic Rain Forest *sensu lato*, where *F. citrifolia* and *F. eximia* are native.

Table 1 - Transferability and characterization of *Ficus insipida* (prefix Fins)¹, *Ficus racemosa* (prefix Frac)² and *Ficus citrifolia* (prefix Frub)² microsatellite loci in 60 *Ficus citrifolia* and 60 *Ficus eximia* individuals from two populations in southeastern Brazil.

Locus accession n.	Primer sequence 5'-3'	Repeat motif	Size range (bp)	Characterization of microsatellite markers									
				<i>Ficus citrifolia</i>					<i>Ficus eximia</i>				
				bp _o	N _a	T _a	H _o /h _E	Pe	bp _o	N _a	T _a	H _o /h _E	Pe
FinsN1	F: AGGGCTGAGATAGGTTGATT R: TAAAGTTGGTGTGGCATC	(TA) ₂ (CA) ₁₀ (TA) ₇	150-160	4	50	0.93/0.68-	0.248	158-164	4	49	0.93/0.66-	0.231	
AM039805	F: GAATCTGGAGGTGGAATAAAC R: AAAGATCGCTCGTCAACC	(TA) ₁₁ (TG) ₁₆	193-210	5	46	0.17/0.31-	0.051	178	1	50	*	-	
FinsU9	F: CGTGTATTGATGTGTGTG R: TCACCTCTCTCTCTTTTG	(AG) ₁₆	148-156	-	-	-	-	-	-	-	-	-	
AM039811	F: TGTCACCTGTTCTGTTTGTG R: CAGCCAAACCCCAAGTATAAGA	(TC) ₁₃ (CA) ₁₀	153-167	15	46	0.90/0.91	0.662	162-178	8	50	0.25/0.83-	0.466	
Frac86	F: CCAGAACAGGTTGGACGTAAC R: GGATTACCCGGCTATGAAAT	(CA) ₁₃	166-172	-	-	-	-	-	-	-	-	-	
DQ659281	F: ACCAAGAGCCAAACTCGT R: TCAACCTTGTGCTCTTTC	(AC) ₁₃	147-151	1	48	*	-	144-159	6	48	1.00/0.78-	0.380	
Frac110	F: CCACCTTGAATGTCACCTGGA R: TGAACAGCCAACTGAGAATG	(AG) ₂₄	188-226	8	48	0.73/0.78	0.399	195-255	12	50	0.72/0.87-	0.577	
DQ659282	F: ACACGTGCAGTGTGCTGA R: ACAGCTGCCCAATTCCTTGA	(AG) ₈ AAAG(GA) ₁₃	195-255	6	50	0.67/0.77	0.359	192-222	7	49	0.36/0.78-	0.383	
Frac154	F: GTACACTCTTAAAGCTGCC R: GTACACTCTTAAAGCTGCC	(TC) ₂₄	145-188	12	50	0.55/0.82-	0.468	120-165	11	50	0.64/0.87-	0.561	
DQ659284	F: TAITTCAAATAACATCTCTCAAC R: TACGTTTGTATGGACTTTGGC	(GA) ₁₁	106-136	-	-	-	-	-	-	-	-	-	
Fub29	F: AGATGTCAAAATAAGGTCAGCT R: AGATGCAGTTCATACAAATC	(TG) ₁₉	149-173	9	49	0.46/0.83-	0.485	140-156	5	50	0.95/0.77-	0.352	
DQ659290	F: GCACGTAGTCGGTGTAAAGC R: CTGTGCGGAATAAAAAGCTAGC	(AC) ₁₀	150-173	4	50	0.58/0.54	0.141	158-164	2	48	0.39/0.42	0.087	
Fub38	F: CAGCAATGATCTTGACCT R: ACTCATCAATACTCTAAACAAC	(CA) ₁₄ (CA) ₈	228-246	6	49	1.00/0.79-	0.391	215-230	4	49	0.93/0.64-	0.216	
DQ659291	F: GCGTGAAAATAIGCTATGA R: GTTTCGTTCAAATTTGAGTGC	(AC) ₁₈	172-190	8	49	0.48/0.80-	0.426	156-170	8	49	0.70/0.79	0.407	
Fub391	F: GTACTGTGATTAGTATCTTTGA R: CTAGCAATAAATCACTGATATTG	(AC) ₂₂	135-159	9	48	0.55/0.83-	0.478	148-188	9	48	0.53/0.82-	0.462	
DQ659292	Sum ¹ /Average ² /Cumulative probability of exclusion ³		87/17.3 ²			0.58 ² /0.67 ²	0.996 ³	77/16.4 ²			0.62 ² /0.69 ²	0.995 ³	

1, Vignes *et al.* 2006; 2, Crozier *et al.* 2007; size range (bp), original data for species for which primers were developed; bp_o, observed size range; N_a, number of alleles; T_a, annealing temperature (°C); H_o, observed heterozygosity; H_E, expected heterozygosity; Pe, paternity exclusion probability; *, FinsT7 and Frac154 were monomorphic in just one of the species; -, statistically significant deviation from Hardy-Weinberg equilibrium (p < 0.05).

Acknowledgments

We thank the Instituto Florestal for permitting this research (ref. # 43.487/2007) and for the use of the facilities in protected areas. This study was supported by grants from FAPESP (R.A.S.P #04/10299-4, A.L.A.M. #03/04199-4/2004) the Provost for Research of Sao Paulo University (A.L.A.M.) and FAEPA (M.A.M.). A.G.N. was supported by a fellowship from CAPES and J.M.F. from FAPESP (#2006/05152-0). The authors thank Marcela Corbo Guidugli for technical support on microsatellite analyses and the anonymous referees for their useful comments on a previous version of the manuscript.

References

- Ahmed S, Dawson DA, Compton SG and Gilmartin PM (2007) Characterization of microsatellite loci in the African fig *Ficus sycomorus* L. (Moraceae). *Mol Ecol Notes* 7:1175-1177.
- Bandelj D, Javornik B and Jakse J (2007) Development of microsatellite markers in the common fig, *Ficus carica* L. *Mol Ecol Notes* 7:1311-1314.
- Berg CC (1989) Classification and distribution of *Ficus*. *Experientia* 45:605-611.
- Crozier YC, Jia XC, Yao JY, Field AR and Cook JM (2007) Microsatellite primers for *Ficus racemosa* and *Ficus rubiginosa*. *Mol Ecol Notes* 7:57-59.
- Dick CW, Hardy OJ, Jones FA and Petit RJ (2008) Spatial scales of pollen and seed-mediated gene flow in tropical rain forest trees. *Trop Plant Biol* 1:20-33.
- Doyle JJ and Doyle JL (1990) Isolation of plant DNA from fresh tissue. *Focus* 12:13-15.
- Epperson BK and Alvarez-Buylla ER (1997) Limited seed dispersal and genetic structure in life stages of *Cecropia obtusifolia*. *Evolution* 51:275-282.
- Giraldo E, Viruel MA, López-Corrales M and Hormaza JI (2005) Characterization and cross-species transferability of microsatellites in the common fig (*Ficus carica* L.). *J Horticult Sci Biotechnol* 80:217-224.
- Goudet J (2001) FSTAT, v. 1.2: A computer program to calculate F-statistics. *Heredity* 86:485-486.
- Hamrick JL and Loveless MD (1989) The genetic structure of tropical tree populations: Associations with reproductive biology. In: Bock JL and Linhart YB (eds) *The Evolutionary Ecology of Plants*. Westview Press, Boulder, pp 129-146.
- Kalinowski ST, Taper ML and Marshall TC (2007) Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Mol Ecol* 16:1099-1106.
- Khadari B, Hochu I, Santoni S and Kjellberg F (2001) Identification and characterization of microsatellite loci in the common fig (*Ficus carica* L.) and representative species of the genus *Ficus*. *Mol Ecol Notes* 1:191-193.
- Kim K-S, Min M-S, An J-H and Lee H (2004) Cross-species amplification of Bovidae microsatellites and low diversity of the endangered Korean Goral. *J Hered* 95:521-525.
- Kinnaird MF, O'Brien TG and Suryadi S (1996) Population fluctuation in Sulawesi Red-Knobbed Hornbills: Tracking figs in space and time. *The Auk* 113:431-440.
- Moon HS, Nicholson JS and Lewis RS (2008) Use of transferable *Nicotiana tabacum* L. microsatellite markers for investigating genetic diversity in the genus *Nicotiana*. *Genome* 51:547-599.
- Nason JD, Herre EA and Hamrick JL (1996) Paternity analysis of the breeding structure of strangler fig populations: Evidence for substantial long-distance wasp dispersal. *J Biogeogr* 23:501-512.
- Nazareno AG and Carvalho D (2008) What the reasons for no inbreeding and high genetic diversity of the neotropical fig tree *Ficus arpazusa*? *Conserv Genet* (DOI 10.1007/s10592-008-9776-x).
- Poncet V, Hamon P, Minier J, Carasco C, Hamon S and Noirot M (2004) SSR cross-amplification and variation within coffee trees (*Coffea* spp.). *Genome* 47:1071-1081.
- Powell W, Machray GC and Provan J (1996) Polymorphism revealed by simple sequence repeats. *Trends Plant Sci* 1:215-222.
- Sanguinetti CJ, Dias EN and Simpson AJG (1994) Rapid silver staining and recovery of PCR products separated on polyacrylamide gels. *Biotechnology* 17:914-921.
- Shanahan M, Compton SG, So S and Corlett R (2001) Fig-eating by vertebrate frugivores: A global review. *Biol Rev* 76:529-572.
- Vignes H, Hossaert-McKey M, Beaune D, Fevre D, Anstett MC, Borges RM, Kjellberg F and Chevallier MH (2006) Development and characterization of microsatellite markers for a monoecious *Ficus* species, *Ficus insipida*, and cross-species amplification among different sections of *Ficus*. *Mol Ecol Notes* 6:792-795.
- Weiblen GD (2002) How to be a fig wasp. *Annu Rev Entomol* 47:299-330.
- Zavodna M, Arens P, Van Dijk PJ and Vosman B (2005) Development and characterization of microsatellite markers for two dioecious *Ficus* species. *Mol Ecol Notes* 5:355-357.

Associate Editor: Márcio de Castro Silva Filho

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.